



E-ISSN: 2788-8428
P-ISSN: 2788-8436
IJZEL 2022; 2(1): 81-85
Received: 15-02-2022
Accepted: 20-03-2022

Mark Ian Cooper
University of Johannesburg,
Auckland Park 2006, Gauteng,
South Africa

Is mass correlated with width among red millipedes *Centrobolus* Cook, 1897?

Mark Ian Cooper

Abstract

Body mass variation was correlated with male and female width in the red millipede genus *Centrobolus*. Mass and width ($r=0.91$, Z score= 3.46 , $n=8$, $p<0.01$) in males ($r=0.91$, Z score= 2.13 , $n=5$, $p=0.02$) and females ($r=0.995$, Z score= 4.27 , $n=5$, $p<0.01$) were correlated. *C. inscriptus* females have the largest width (7.02 mm) and the highest mass (2.61 g) while *C. digrammus* has the lowest width (4.00 mm) and the lightest mass (0.68 g).

Keywords: Dimorphic, female, mass, morphology, ratio, width

1. Introduction

Red millipedes are found in the southern African subregion with northern limits on the east coast being about -17° latitude S and southern limits being -35° latitude S. They are well represented in the littoral forests of the eastern half of the subcontinent [28]. It consists of taxonomically important species with 12 species considered threatened and includes nine vulnerable and three endangered species [30]. It occurs in all the forests of the coastal belt from the Cape Peninsula to Beira in Mocambique [29]. In an analysis of four morphometric factors (male and female length and width), four positive correlations were found in the 22 species of *Centrobolus* [25]. These worm-like millipedes have female-biased sexual size dimorphism [7-25]. Here, body mass is correlated with body width in *Centrobolus* Cook, 1897 [6, 29, 30]. One hypothesis is body width correlates with body mass. The aim of the study is to determine the exact relationship between mass and width through correlation analysis with different species.

2. Materials and Methods

C. digrammus, *C. fulgidus*, *C. inscriptus*, *C. ruber* were identified as belonging to the genus *Centrobolus* Cook, 1897 from field collections at Admirals Waterfall, Richard's Bay, Mtunzini, and Anerley; all in South Africa (Table 1). Millipede collection localities were made in the rainy season [7]. Weight in millipedes was calculated with a (Mettler AC 100) Autobalance. Body width was calculated as the horizontal body width using a pair of vernier calipers calibrated in millimeters accurate to three significant figures. Horizontal tergite width and mean body mass were correlated with a Pearson Correlation Coefficient calculator available at <https://www.gigacalculator.com/calculators/correlation-coefficient-calculator.php>.

3. Results

Mass and width (Fig. 1: $r=0.93468004$, Z score= 4.48243126 , $n=10$, $p=0.00000369$) in males (Fig. 2: $r=0.90615676$, Z score= 2.12925309 , $n=5$, $p=0.01661660$) and females (Fig. 3: $r=0.99521810$, Z score= 4.26644915 , $n=5$, $p=0.00000994$) were related. Body width was normally distributed ($D=0.18655$, $n=10$, $p=0.81643$). Body mass was normally distributed ($D=0.23289$, $n=10$, $p=0.57311$). Female width was normally distributed ($D=26643$, $n=5$, $p=0.79033$). Female mass was normally distributed ($D=0.29699$, $n=5$, $p=0.67584$). Male width was normally distributed ($D=21396$, $n=5$, $p=0.93695$).

Correspondence

Mark Ian Cooper
University of Johannesburg,
Auckland Park 2006, Gauteng,
South Africa

Table 1: Species in the millipede genus *Centrobolus* Cook, 1897, with body width and body mass. Male followed by female body masses are given with sample sizes in parentheses. Two sets of measurements are included for *C. inscriptus*

Species	width	Location	Body mass (g)
<i>C. fulgidus</i>	5.4 (11), 6.2 (11)	Richards Bay	1.29 (11), 1.97 (11)
<i>C. inscriptus</i>	5.9 (88), 5.98 (56), 6.7 (88), 7.02 (41)	Mtunzini	2.48 (88), 2.00 (56); 2.27 (88), 2.61 (41)
<i>C. ruber</i>	5.0 (18), 6.1 (18)	Port Shepstone	1.28 (18), 2.00 (18)
<i>C. digrammus</i>	4.0 (6), 4.8 (6)	Admirals Waterfall	0.68 (6), 1.02 (6)




Fig 1: Relationship between body width (x-axis) and body mass (y-axis) in *Centrobolus* Cook, 1897




Fig 2: Relationship between male body width (x-axis) and male body mass (y-axis) in *Centrobolus* Cook, 1897




Fig 3: Relationship between female body width (x-axis) and female body mass (y-axis) in *Centrobolus* Cook, 1897

4. Discussion

Significant positive relationships between mass and length and width are given. *C. inscriptus* females have the largest width (7.02 mm) and the highest mass (2.61 g) while *C. digrammus* has the lowest width (4.00 mm) and the lightest mass (0.68 g). This study supports body mass and body width as predictable in *Centrobolus*.

Size-assortative mating based on body mass and body width determined through the variance in millipede polygynandrous mating systems across a size gradient with lower body width occurring in lighter individuals is shown. Body width increases with heavier body mass which is explained by a combination of copulatory guarding and fecundity selection.

In Danish red foxes *Vulpes vulpes* SSD varies with body mass at different population densities ^[31]. In the wandering albatross *Diomedea exulans* SSD varies with body mass with different adult survival, breeding probability, breeding success, chick mass, and juvenile survival ^[26]. In microtines mass rather than length is used to measure SSD ^[5]. In five species of *Drosophila*, the degree of dimorphism in development time was significantly correlated with dry weight and fecundity, with lighter species tending to be more dimorphic for development time as well as more fecund, both in absolute terms and terms of fecundity per unit weight ^[2]. In Red-Eared Sliders *Trachemys scripta elegans* sexual dimorphism of size in favor of females is maintained for variables related to mass and size ^[27]. Millipedes resemble the patterns of mass and sexual dimorphism in adult dragonflies ^[1]. Sexual dimorphism in mass and protein content of the forelimb muscles is known in the northern leopard frog *Rana pipiens* ^[3]. The influence of mass and mating systems has been documented in the most dimorphic mammals ^[32]. Morphological variance in the polygynandrous reproductive system of male and female millipedes means lighter individuals have lower body width or heavier individuals have greater body width pointing toward the evolution of condition-dependent SSD ^[4].

5. Conclusion

The width varied systematically with mass in *Centrobolus*. Morphological variance in the polygynandrous reproductive systems of *Centrobolus* has lighter individuals with a narrower width, or conversely heavier individuals have greater body width. Mass and width impact the moments of inertia.

6. Competing interests

The author has declared that no competing interests exist.

7. References

- Anholt B, Marden J, Jenkins D. Patterns of Mass Gain and Sexual Dimorphism in Adult Dragonflies (Insecta, Odonata). *Canadian Journal of Zoology*. 2011;69(5):1156-1163.
- Bharathi NS, Prasad NG, Shakarad M, Joshi A. Correlates of sexual dimorphism for dry weight and development time in five species of *Drosophila*. *Journal of Zoology*. 2004;264(1):87-95.
- Blackburn D. Sexual dimorphism in mass and protein content of the forelimb muscles of the northern leopard frog, *Rana pipiens*. *Canadian Journal of Zoology*. 1992;70(4):670-674.
- Bonduriansky R. The Evolution of Condition-Dependent Sexual Dimorphism. *The American Naturalist*. 2007; 169(1): 9-19.
- Boonstra R, Gilbert BS, Krebs CJ. R. Mating Systems and Sexual Dimorphism in Mass in Microtines. *Journal of Mammalogy*. 1993;74(1):224-229.
- Cook OF. New relatives of *Spirobolus giganteus*. *Brandtia* (A series of occasional papers on Diplopoda and other Arthropoda). 1897;18(2):73-75.
- Cooper MI. Mating dynamics of South African forest millipedes *Centrobolus* (Diplopoda: Pachybolidae). The University of Cape Town, 1998, 1-141.
- Cooper MI. Sexual size dimorphism and corroboration of Rensch's rule in *Chersastus* millipedes. *Journal of Entomology and Zoology Studies*. 2014;2(6):264-266.
- Cooper MI. Copulation and sexual size dimorphism in worm like millipedes. *Journal of Entomology and Zoology Studies*. 2017;5(3):1264-1266.
- Cooper M. *Centrobolus anulatus* (Attems, 1934) reversed sexual size dimorphism. *Journal of Entomology and Zoology Studies*. 2018;6(4):1569-1572.
- Cooper MI. The relative sexual size dimorphism of *Centrobolus inscriptus* compared to 18 congeners. *Journal of Entomology and Zoology Studies*. 2016;4(6):504-505.
- Cooper MI. Relative sexual size dimorphism in *Centrobolus digrammus* (Pocock) compared to 18 congeners. *Journal of Entomology and Zoology Studies*. 2017;5(2):1558-1560.
- Cooper MI. Relative sexual size dimorphism in *Centrobolus fulgidus* (Lawrence) compared to 18 congeners. *Journal of Entomology and Zoology Studies*. 2017;5(3):77-79.
- Cooper MI. Relative sexual size dimorphism *Centrobolus ruber* (Attems) compared to 18 congeners. *Journal of Entomology and Zoology Studies*. 2017;5(3):180-182.
- Cooper MI. Competition affected by re-mating interval in a myriapod. *Journal of Entomology and Zoology Studies*, 3(4), 77-78.
- Cooper M. Re-assessment of Rensch's rule in *Centrobolus*. *Journal of Entomology and Zoology Studies*. 2017;5(6):2408-2410.
- Cooper MI. Sexual size dimorphism and the rejection of Rensch's rule in Diplopoda. *Journal of Entomology and Zoology Studies*. 2018;6(1):1582-1587.
- Cooper MI. Allometry for sexual dimorphism in millipedes. *Journal of Entomology and Zoology Studies*. 2018;6(1):91-96.
- Cooper MI. Trigoniulid size dimorphism breaks Rensch. *Journal of Entomology and Zoology Studies*. 6(3), 1232-1234.
- Cooper M. A review of studies on the fire millipede genus *Centrobolus* (Diplopoda: Trigoniulidae). *Journal of Entomology and Zoology Studies*. 2018;6(4):126-129.
- Cooper M. *Centrobolus sagatinus* sexual size dimorphism based on differences in horizontal tergite widths. *Journal of Entomology and Zoology Studies*. 2018;6(6):275-277.
- Cooper M. *Centrobolus silvanus* dimorphism based on tergite width. *Global Journal of Zoology*. 2018;3(1):003-005.

23. Cooper M. Xylophagous millipede surface area to volume ratios are size dependent in forest. *Arthropods*. 2019;8(4):127-136.
24. Cooper M. Mass covaries with volume in forest millipedes *Centrobolus* Cook, 1897. *Journal of Entomology and Zoology Studies*. 2021;9(6):190-192.
25. Cooper M. Length and Width Correlations in *Centrobolus* Cook, 1897. *New Visions in Biological Science*. 2022;9:39-45.
26. Cornioley T, Jenouvrier S, Borger L, Weimerskirch H, Ozgul A. Fathers matter: male body mass affects life-history traits in a size-dimorphic seabird. *Proceedings of the Royal Society B*. 2017;284(1854):20170397.
27. Gradela A, Santiago TOC, Pires IC, de Castro Souza Silva A, de Souza LC, de Faria MD, *et al*. Sexual Dimorphism in Red-Eared Sliders (*Trachemys scripta elegans*) from the Wild Animal Triage Center of the Tiete Ecological Park, São Paulo, Brazil. *Acta Scientiae Veterinariae*. 2017;45(1468):1-10.
28. Hamer ML. Checklist of Southern African millipedes (Myriapoda: Diplopoda). *Annals of the Natal Museum*. 1998;39(1):11-82.
29. Lawrence RF. The Spiroboloidea (Diplopoda) of the eastern half of Southern Africa*. *Annals of the Natal Museum*. 1967;18(3):607-646.
30. Mailula RP. Taxonomic revision and Red List assessment of the red millipede genus *Centrobolus* (Spirobolida: Pachybolidae) of South Africa. University of Kwazulu natal, 2021, 23+289.
31. Pagh S, Hansen MS, Jensen B, Pertoldi C, Chriél M. Variability in body mass and sexual dimorphism in Danish red foxes (*Vulpes vulpes*) in relation to population density. *Zoology and Ecology*. 2018;28(1):1-9.
32. Weckerly FW. Sexual-Size Dimorphism: Influence of Mass and Mating Systems in the Most Dimorphic Mammals. *Journal of Mammalogy*. 1998;79(1):33-52.