



E-ISSN: 2788-8428
P-ISSN: 2788-8436
ZEL 2023; 3(1): 32-36
Received: 01-12-2022
Accepted: 05-01-2023

Biju Adhikari
Faculty of Agriculture,
Agriculture and Forestry
University, Rampur, Chitwan,
Nepal

Santosh Khadka
Department of Entomology,
Agriculture and Forestry
University, Rampur, Chitwan,
Chitwan, Nepal

Corresponding Author:
Biju Adhikari
Faculty of Agriculture,
Agriculture and Forestry
University, Rampur, Chitwan,
Nepal

Zoological and Entomological Letters

Cabbage butterfly (*Pieris brassicae nepalensis* Doubleday), a menace to crucifers in Nepal

Biju Adhikari and Santosh Khadka

Abstract

Crucifers are significant commercially grown vegetable crops that are produced throughout the world, especially in Nepal's high highlands and terai areas. Numerous significant insect pests usually attack them. The cabbage butterfly is one of them and is a worldwide nuisance of cruciferous vegetables like cabbage, cauliflower, and Knol Khol. It can also attack radish, turnip, Sarson, toria and other cruciferous plants. In the Terai and mid hills, the subspecies *Pieris brassicae nepalensis* outnumbers other subspecies. Early instar larvae munch haphazardly on the leaf, scrapping the leaf lamina before eating holes in it. This results in severe crop loss, and in the worst cases, crop collapse. In Nepal, the late-season crops of cabbage and cauliflower are severely harmed by the cabbage butterfly (*Pieris brassicae nepalensis*), which can result in yearly output losses of up to 80–100%. The cabbage butterfly of cruciferous vegetables is a significant insect pest. This page concentrates on its identification, distribution, host range, life cycle, nature of the damage, and long-term treatment, including IPM measures.

Keywords: Crucifers, cabbage, cabbage butterfly, *Pieris brassicae nepalensis*

1. Introduction

Insect pests are the most important crop-limiting factor for crucifers (Bhavani *et al.*, 2009)^[9]. Each year, they can account for up to 40% of crop loss (Ali & Rizvi, 2007; Hasan & Ansari, 2010; Sood & Bhalla, 1996)^[2, 37, 27]. In Nepal, four cabbage butterfly species have been identified. The species are *P. brassicae*, *P. brassicae nepalensis*, *P. canidida*, and *P. canidida indica* (Thapa, 1987)^[30]. In Nepal's Terai and inner Terai regions, *P. brassicae nepalensis* predominates, followed by *P. candida* (Thapa, 1987)^[30]. *P. brassicae* readily consumes members of five major plant families, including the Cruciferae), Tropaeolaceae, Capparaceae, Resedaceae, and Papilionaceae (Feltwell, 1982)^[12]. *P. brassicae nepalensis*, on the other hand, prefers the Cruciferae, of which cabbage is a major host (Feltwell, 1982)^[12]. *Pieris brassicae nepalensis* Doubleday (Lepidoptera: Pieridae) is a cruciferous plant pest that feeds on cabbage, cauliflower, mustard, turnip, and other plants. It is a serious pest of nearly all crucifers, attacking crops at various stages and causing significant damage and yield losses. They are usually known by their names cabbage worm, big white butterflies, cabbage white, or cabbage moth, and are the most common crucifer lepidopteran pest (Ansari *et al.*, 2012)^[3]. They were given the nickname "cabbage butterfly" because of their snow-white wing and the reality that cabbage is their primary food source (Wilbur, 2011)^[31]. In Nepal, it is a damaging pest that lowers the annual output of late-planted cabbage and cauliflower reaching up to 80–100%. (Joshi, 1994)^[18]. In order to maintain this pest population underneath the level that generates economic impact, integrated pest management (IPM) control is critical.

2. Materials and Methods

This review's content is solely based on secondary sources. Numerous academic journals, agriculture institutions, and some other resources including MoALD, CABI, and pertinent reports were used to construct a sample of the information that is presently accessible. This analysis was then concluded, and the main findings were summarized. The investigation also took advice from officials and experts in related disciplines.

3. Results and Discussion

3.1. Taxonomic position

Cabbage butterflies are the most common members of the genus *Pieris*, which belongs to the order Lepidoptera, family Pieridae, and sub-species *nepalensis*.

Kingdom: Animalia
 Phylum: Arthropoda
 Class: Insecta
 Order: Lepidoptera
 Family: Pieridae
 Genus: Pieris
 Species: brassicae
 Subspecies: nepalensis

3.2. Distribution

P. brassicae was discovered in South Africa's Western Cape in 1994 (Claassens, 1995) ^[10]. In February 1996, the first record of *P. brassicae* from Delhi was made. Given that it is primarily a nuisance in mountainous areas, its high prevalence was surprising (Bhalla *et al.*, 1997) ^[7]. This species is found as subspecies *P. brassicae nepalensis* in Nepal, India, Tibet (Xizhang), and Yunnan, China.

3.3. Host range and crop losses

This oligophagous pest feeds on a variety of family members and prefers cruciferous crops. Cabbage and cauliflower are the most commonly attacked host plants, but it also attacks knoll khol, turnip, radish, sarsoo, mustard, and other cruciferous crops. *P. brassicae* (L.) is one of the most destructive and cosmopolitan pests of cruciferous crops in India, accounting for 40% of crop damage each year, according to Hasan and Ansari (2011) ^[14]. Damage to cabbage heads is estimated to be 70 to 98 percent (Prasad, 1963). *P. brassicae* larvae attacked 68.5 percent of the marketable yield of cabbage cv. Pride of India conducted field research in Upper Shillong, Meghalaya, India, in 1991-92. (Thakur, 1996) ^[29]. In 1985 and 1986, *P. brassicae* caused 40.45% damage to cabbages in Izmir, Turkey, and 27.06 percent damage to cauliflowers (Atalay and Hincal, 1992) ^[5]. Crop attacks are more localized and can result in complete crop loss in one area. The pest causes severe damage to crucifer crops at every stage of development, from seedling to flowering (Bhandari *et al.*, 2009) ^[8]. This pest causes losses ranging from 27 to 41% in various conditions (Ali & Rizvi, 2007; Atalay & Hincal, 1992; Eichler, 1948; Shapiro, 1975) ^[2, 5, 11, 25]. Furthermore, because adults tend to migrate, they may infest previously uninfested areas. As a result, because of its ability to spread to new areas, this pest is becoming a threat to our country.

3.4. Seasonal abundance

Research on the population dynamics of the cabbage butterfly indicates that this pest first appears on cruciferous crops in early October and is active through the end of April. From May to September, this pest is scarce from the plain; however, breeding normally occurs in the mountainous area. High temperatures, extended daylight hours, low relative humidity, and little rain all contribute to population increase (Sood & Bhalla, 1996) ^[27]. Abiotic variables can be employed to manage the cabbage butterfly outbreak and stop future damage. *P. brassicae* reproduces three times each year in Nepal, for each generation lasting 32–64 days. During February and May, the first generation wreak havoc on cauliflower and cabbage.

3.5. Life cycle

The life span of *Pieris brassicae* changes based on the surrounding environment. Its life cycle lasts 15 to 22 days from March to April, and 30 to 40 days from November to February. Such weather factors are typical in Nepal's hills and plains, except for the colder months.

3.5.1. Eggs

Under the leaf sheath, the adult moth lays 160–170 eggs on average. Eggs are normally adhered to the backside of the surface of the leaf and are always placed in bunches. Before hatching, the newly produced eggs are yellow in color with grayish-black tips. Eggs have a form resembling a bullet or a maize cob with longitudinal ridges (Kumar *et al.*, 2020) ^[20]. For two to three days, the females continue to deposit eggs in groups beneath the surface of cabbage leaves. The incubation period is influenced by seasonal variations. Eggs are incubated for 3-6 days on average. The incubation period is influenced by seasonal variations.

3.5.2. Larva


There are 5 larvae instars and four molts in the *P. brassicae nepalensis* caterpillar featuring a slightly dark head and a hairy body the newly born caterpillars are cylindrical in shape mild yellow in color and have a smooth exterior. The 3rd instar larvae are often more active and have a greenish body color with black spots during this instar it is typical to observe the larvae devouring their host plant ferociously as the larvae move through the 4th instar they resemble those in the 3rd instar but are bigger and feed differently there is a lot of hair on the 5th instar larvae after reaching this stage they are elongated robust and golden in hue with vivid coloration on their abdomen and thorax (Molet, 2011) ^[22]. Their heads are depicted as being black and grey. To complete development during this instar, the larva needs the best quality and amount of food; otherwise, it will perish before maturing into a mature butterfly (Metspalu *et al.*, 2003) ^[35]. The pupal phase lasts 8–15 days (about two weeks) in March–April and 20–28 days in November–February (about 4 weeks).

3.5.3. Pupa

Under the leaves, pupation takes place. The pupae contain multiple dots and are pale to greyish in hue. The pupae had lateral ridges that ran down both sides, and a dorsal ridge that extended from the centre of the anterior across the entire body. The pupal stage of a butterfly's life cycle typically lasts 7 to 10 days (Hasan *et al.*, 2008) ^[16].

3.5.4. Adult

The cabbage caterpillar's adults are pale white in color. In both males and females, the wings are white with dark brown tips on the forewings. Females have one pair of brownish-black spots on the dorsal side of each forewing, whereas males have none. A female had two black circular spots on the dorsal side of each forewing (Atwal and Dhaliwal, 2015) ^[6]. Adult butterflies have a lifespan of 2.5-12.5 days.



Source: CABI

Fig 1: Biology of cabbage butterfly

3.6. Nature of damage

All of the plant's above-ground components are infested by *Pieris brassicae* at every stage of growth. Caterpillars harm plants by eating in large groups and poking round holes, typically on the outside leaves, in the foliage (Younas *et al.*, 2004; Khalid, 2006) [36, 19]. In contrast to later instar caterpillars, which consume leaves from the periphery inward while leaving the primary veins intact, the very first instar caterpillar just scrapes the leaf surface. If the vegetable's leaves are punctured, it can become unfit for consumption. The caterpillar causes the most damage when it burrows into the crown of cauliflower and cabbage. In extreme infestations, plants succumb and the leaves are completely destroyed. The pest is extremely terrible in the months of April and March.

All development stages of crucifer crops, from juvenile to blossoming, are severely harmed by the insect (Bhandari *et al.*, 2009) [8]. Damage from this pests vary from 27 to 41% depending on the situation. (Eichler, 1948; Shapiro, 1975; Atalay & Hincal, 1992; Ali & Rizvi, 2007; Atalay & Rizvi) [11, 25, 5, 2].



Source: Adhikari

Fig 2: Damage symptoms by cabbage butterfly

3.7. Integrated pest management

3.7.1. Cultural control

In Nepal, uprooting and removing all weeds and previous crop remnants, burning or composting, and intercropping with repellent crops, such as onion and garlic. Plant one

repellent plant for every ten cauliflowers, using the recommended crop spacing, e.g., Kathmandu local = 60X60cm (it's important to avoid dense planting), and the recommended fertilizer dose, with a split dose of nitrogen. 4.5 kg Urea+13 kg DAP +3.5 kg MOP/ropani as a starting dose, then one month later after seedling transplant Some of the cultural controls used to prevent cabbage butterfly infestation include 5.5 kg urea/ropani and routine irrigation to keep the field from becoming too dry (Adhikari *et al.*, 2014) [1]. Planting resistant cultivars, such as the Copenhagen market, also aids in avoiding an infestation. In a Nepalese experiment, the lowest cabbage butterfly population was recorded in the Copenhagen market (10.06 larva/plant).

3.7.2. Mechanical control

In several experiments conducted in 1987 and 1988, cauliflower fields planted from May to July in Germany were covered with netting. Covering the crop for 4-5 weeks after planting was sufficient to exclude and control *P. brassicae* in the first few weeks after planting (Wonneberger and Gawehn, 1989) [32]. Hand-picking leaves with egg masses and larvae, crushing or keeping them in a bucket of soapy water, and collecting and destroying adult butterflies using a sweep net during the day when adults are active have all been used as mechanical control by Nepalese farmers (Adhikari *et al.*, 2014) [1].

3.7.3. Biological control

Apanteles wasps attacked the larvae, laying eggs inside their bodies and spinning cocoons on or near them. Kumar *et al.*, 2020 discovered that Apanteles sp. parasitized larvae exhibited no visible symptoms until the apanteles cocoon formed or the parasitoid emerged. Apanteles parasitizes *P. brassicae* larvae in their natural habitat (Atwal and Dhaliwal, 2015) [6]. The larval survival rate of *P. brassicae* has been reported to be 83%. 1999 (Aslam and Suleman). In Nepal, foliar application of Azadirachtin (neem-based insecticide) at 5 ml /lit water and 25 lit water /ropani; maximum three sprays, at one-week intervals, and foliar application of Beauveria bassiana (e.g.'Daman') 1% WP @ 5 gm /lit water and 25 lit water /ropani during the late afternoon (to avoid the sun); maximum three sprays (Adhikari *et al.*, 2014) [1]. According to Jaquet *et al.*, *P.*

brassicae larvae were highly susceptible to purified crystals of *Bacillus thuringiensis* (Bt) subsp. *thuringiensis* (1986)^[17]. *Metarhizium anisopilae* and *Beauveria bassiana* were also effective against the pest. CAMB Bt-based biopesticides were found to be effective against *P. brassicae* on cauliflower (Zafar *et al.*, 2002)^[34]. Spinosad and Cypermethrin, for example, were more effective in causing death and reducing leaf damage (Giri *et al.*, 2020)^[13].

3.7.4. Chemical control

In field trials in Manipur, India, Ram and Pathak (1992)^[24] found that carbaryl and dimethoate were effective at reducing *P. brassicae* infestation on cabbage. Eleven insecticides were tested for *P. brassicae* control in cabbage cv. Pride of India grown in Barapani, Meghalaya, India, in 1991-92. The plants were sprayed twice, once before and once after heading. Overall, fenvalerate was almost completely effective against *P. brassicae*, followed by deltamethrin (97.3%), cypermethrin (96.8%), malathion (96.08%), and fenitrothion (93.3%). Chlorpyrifos, quinalphos, and diflubenzuron were the least effective insecticides tested, while fenvalerate-treated plots yielded the highest yield.

Ram and Pathak (1992)^[24] reported that carbaryl and dimethoate were effective in reducing *P. brassicae* infestation on cabbage in field trials in Manipur, India. During 1991-92, eleven insecticides were tested for control of *P. brassicae* in cabbage cv. Pride of India grown in Barapani, Meghalaya, India. Plants were sprayed twice, once before heading and again after heading. Overall, fenvalerate controlled *P. brassicae* almost completely, followed by deltamethrin (97.3%), cypermethrin (96.8%), malathion (96.08%), and fenitrothion (93.3%). The least effective insecticides tested were chlorpyrifos, quinalphos, and diflubenzuron whereas the highest yield was obtained in fenvalerate-treated plots. Fenvalerate also had the highest cost-benefit ratio (Thakur and Deka, 1997)^[28]. Foliar application of Malathion (50 EC) at a rate of 2 ml /lit water and 25 lit water /ropani, Emamectin Benzoate (5% SG) at a rate of 2 gm /16 lit water and 25 lit water /ropani, and Cypermethrin (10 EC) at a rate of 2 ml /lit water and 25 lit water /ropani, and Cypermethrin (Adhikari *et al.*, 2014)^[1].

4. Conclusion

The oligophagous cabbage butterfly is a pest that consumes cruciferous plants. While attacking Brussels sprouts, mustard, and other vegetables, it favors cabbage and cauliflower. To control this destructive crucifer pest, IPM techniques like cultural control, mechanical control, biological control, and chemical control may be used. The right date of planting may aid in preventing pest incidence by severing the link between crop stage and pest appearance. In order to decrease the number of adults, hand picking and netting are useful mechanical IPM tools. For keeping an eye on pests in new places, it is a very useful technique. In cultural techniques, companion planting with nasturtium, marigold, onion, and garlic is also possible. Resistant/tolerant kinds exceed chemical control as a damaging larval stage. Entomopathogenic fungi like *M. anisopliae* and *B. bassiana* can be utilized to lessen this pest. *Beauveria bassiana*, cow urine solution with water, and foliar spray of the neem-based pesticide Azadirachtin are some biological controls for this pest. This pest was effectively controlled by insecticides like Cypermethrin (10

EC) at 2 ml/lit water and 25 lit water/ropani and Emamectin Benzoate (5% SG) at 2 gm/lit water and 25 lit water/ropani.

5. Acknowledgement

A review paper incorporates data from a wide range of verified sources, therefore we would like to convey our appreciation to all authors for contributing sources cited in this paper.

6. References

1. Adhikari KB, Tripathi N, Dhakal AP. Cabbage butterfly of cauliflower-Nepal. Cabbage butterfly of cauliflower-Nepal. Plantwise Plus Knowledge Bank. CABI International; c2014.
2. Ali A, Rizvi P. Developmental response of cabbage butterfly, *Pieris brassicae* L. (Lepidoptera: Pieridae) on different cole crops under laboratory and field conditions. Asian Journal of Plant Sciences. 2007;6(10):1241-1245.
3. Ansari MS, Hasan F, Ahmad N. Influence of various host plants on the consumption and utilization of food by *Pieris brassicae*. Bulletin of Entomological Research. 2012;102(2):231-237.
4. Aslam M, Suleman N. Biology of *Pieris brassicae* (Linnaeus) (Lepidoptera: Pieridae) under laboratory conditions Pakistan Journal of Biological Sciences. 1999;2(1):199-200.
5. Atalay R, Hincal P. Investigations on the species of Pieridae (Lepidoptera) which are harmful to plants belonging to the family Cruciferae with their importance in Izmir and its vicinity and the biology of the large white butterfly (*Pieris brassicae* L.) along with the factors affecting its population fluctuations. 1992;6(1):271-286.
6. Atwal AS, Dhaliwal GS. Agricultural pests of South Asia and their management. Eighth edition, Kalyani Publishers, Ludhiana; c2015. p. 452.
7. Bhalla S, Kapur ML, Verma BR, Singh R. An unusual abundance of cabbage butterfly, *Pieris brassicae* (Linnaeus) on various brassica species in the environs of Delhi for locating sources of resistance among different cultivars. Journal of Entomological Research. 1997;21(2):147-151.
8. Bhandari K, Sood P, Mehta H, Choudhary A, Prabhakar CS. Effect of botanical extracts on the biological activity of granulosis virus against *Pieris brassicae*. Phytoparasitica. 2009;37(4):317-322.
9. Bhavani, B., Venugopal R, V Ch. Evaluation of different IPM modules for the management of insect pests of Cabbage. Pestology. 2009;33(12):9-13.
10. Claassens AJM. Observations on the large white, *Pieris brassicae* (L.) (Lepidoptera: Pieridae), a butterfly which recently established itself in the Western Cape. Metamorphosis. 1995;6(2):86-93.
11. Eichler D. Failure of cole crops caused by cabbage butterfly. Mitteilungen Der Munchener Entomologischen Gesellschaft. 1948;34(2):417-419.
12. Feltwell J. Large white butterfly, the biology, biochemistry and physiology of *Pieris brassicae* (Linnaeus). The Hague, the Netherlands. Dr. W. Junk Publishers; c1982. p. 535.
13. Giri HN, Sharma MD, Thapa RB, Pande KR, Khatri BB. Efficacy testing of 'soft' pesticides for cabbage butterfly (*Pieris brassicae nepalensis* Doubleday) at

- Rampur, Chitwan. Journal of Agriculture and Forestry University. 2020;4:177.
14. Hasan F, Ansari MS. Effect of different cole crops on the biological parameters of *Pieris brassicae* (L.) (Lepidoptera: Pieridae) under laboratory conditions. Journal of Crop Science and Biotechnology. 2011;13(3):195-202.
 15. Ghode N, Choudhary AS, Raj S, Sahu S, Kumar A. Effect of weed management on yield parameters of cauliflower. Int. J. Agric. Food Sci. 2022;4(1):17-20. DOI: 10.33545/2664844X.2022.v4.i1a.60
 16. Hasan W, Gupta AK, Singh CP. Biology of cabbage butterfly, *Pieris brassicae* (L.) (Lepidoptera: Pieridae) on cabbage and Indian mustard. Journal of Oilseeds Research. 2008;25:104-5.
 17. Jaquet F, Hutter R, Luthy P. Specificity of *Bacillus thuringiensis* Delta-Endotoxin. Applied and Environmental Microbiology. 1986;53(3):500-504.
 18. Joshi SL. Major harmful insect pest of vegetables in Nepal. Vegetable Division, Khumaltar, Nepal; c1994. p. 187.
 19. Khalid S. Bio-ecological studies of *Pieris brassicae* (Lepidoptera: Pieridae) on different hosts. MSc. Thesis, AMU, Aligarh, India; c2006.
 20. Kumar P, Brar JS, Singh G. Life cycle of cabbage caterpillar, *Pieris brassicae* Linn. (Lepidoptera: Pieridae) on cabbage leaves in Talwandi Sabo (Punjab); c2020.
 21. Rajput S, Bisen RK, Raj S, Varma SK, Agrawal HP. Performance of cauliflower varieties under different spacings in Chhattisgarh plain. Int. J. Agric. Food Sci. 2020;2(2):04-07. DOI: 10.33545/2664844X.2020.v2.i2a.34
 22. Molet T. CPHST Pest Datasheet for *Pieris brassicae*. USDA-APHIS-PPQ-CPHST; c2011.
 23. Prasad SK. Quantitative estimation of damage to crucifers caused by cabbage worm, cabbage looper, diamond back moth and cabbage aphid. Indian Journal of Entomology. 1963;25(3):242-259.
 24. Ram S, Pathak KA. Insecticidal control of cabbage butterfly, *Pieris brassicae* Linn. (Pieridae: Lepidoptera) in Manipur. Indian Journal of Entomology. 1992;54(3):353-355.
 25. Shapiro VA. Ecological behavioral aspects of consistence in six crucifers feeding Pierid butterflies in the central Sierra Nevada. Midland Naturalist. 1975;93:424-433
 26. Singh NB, Dhungana S, Adhikari S, Chapagain D, Ghimire N, Sarad DC. Field Screening of Seven Cultivars of Cabbage against Cabbage Butterfly (*Pieris brassicae*) and Cabbage Aphids (*Brevicoryne brassicae*) at Gkuleshor, Baitadi, Nepal. Nepalese Horticulture. 2020;14(1):63-67.
 27. Sood AK, Bhalla OP. Ecological Studies on the Cabbage White Butterfly in the Mid-Hills of Himachal Pradesh. Journal of Insect Science. 1996;9:122-125.
 28. Thakur NSA, Deka TC. Bioefficacy and economics of different insecticides against *Pieris brassicae* (L.) on cabbage in midhills of North-East India. Indian Journal of Plant Protection. 1997;25(2):109-114; 17 ref.
 29. Thakur, NSA. Relationship of cabbage butterfly larval (*Pieris brassicae* L.) population on the marketable yield of cabbage. Journal of Hill Research. 1996;9(2):356-358.
 30. Thapa RB. Biology of *Pieris brassicae nepalensis* Doubleday (Lepidoptera: Pieridae) in the Chitwan Valley. Pesticides. 1987;21(6):30-33.
 31. Wilbur ND. Biological aspects of cabbage white butterfly species, *Pieris brassicae* (Linnaeus, 1758) in the Environs of Taunggyi District, Southern Shan. Universities Research Journal. 2011;4:1-10
 32. Wonneberger C, Gawehn G. Practical results of applying netting to cauliflowers. Gemuse (Munchen). 1989;25(3):164-167.
 33. Teshome Y, Loyew A. Agronomic management status in cauliflower cultural practices in Ethiopia for vegetable crop production technology. Int. J. Agric. Food Sci. 2021;3(1):10-13. DOI: 10.33545/2664844X.2021.v3.i1a.44
 34. Zafar U, Idrees N, Ahmad S, Ahmed AR, Muhammad S, Sheikh R. Performance evaluation of cambiopesticides to control cabbage butterfly (*Pieris brassicae*) in cauliflower crop. Pakistan Journal of Biological Sciences. 2002;5:1041-1043.
 35. Metspalu L, Hiiesaar K, Joudu J, Kuusik A. Influence of Food on Growth, Development and Hibernation of Large White Butterfly. Agronomy Research. 2003;1:85-92.
 36. Younas M, Naeem M, Raqib A, Masud S. Population dynamics of cabbage butterfly (*Pieris brassicae*) and cabbage aphid (*Brevicoryne brassicae*) on five cultivars of cauliflower at Peshawar. Asian Journal of Plant Sciences. 2004;3(3):391-393.
 37. Hasan F, Ansari MS. Effect of different cole crops on the biological parameters of *Pieris brassicae* (L.) (Lepidoptera: Pieridae) under Laboratory Conditions. Journal of Crop Science and Biotechnology. 2010;13:195-202.