

E-ISSN: 2788-8428
P-ISSN: 2788-8436
ZEL 2023; 3(2): 20-29
Received: 02-05-2023
Accepted: 05-06-2023

Dr. Yeshiwas Tarekegn
HOD, Department of Animal
Health, Veterinary
Epidemiologist, Alage ATVET
College, Ziway, Ethiopia

Review on current epidemiological status and public health importance of bovine tuberculosis in Ethiopia

Dr. Yeshiwas Tarekegn

DOI: <https://doi.org/10.22271/letters.2023.v3.i2a.65>

Abstract

The review conducted in these papers revealed the prevalence estimate of *B. tuberculosis* in Ethiopia was determined to be 5.8%. Bovine tuberculosis (BTB) is a known endemic illness of cattle in Ethiopia. The prevalence of intensive dairy production systems was higher (3.1-68%) than that of large livestock production systems (3.4-6.2%) and slaughterhouses (3.5-32%) in different regions of the nation. In Holstein-Friesians, the incidence was higher (14.7-30.7%) than in local zebu (3.4-4.9%) and was linked to genetic resistance. In comparison to the prevalence seen in the established dairy belt in the central areas of Ethiopia, particularly in and around Addis Ababa City, the review also revealed comparatively low average prevalence in the rising dairy districts. Because there are currently fewer animals affected in a smaller geographic area, disease control measures like milk pasteurization, meat inspection, raising awareness about the use of raw milk and milk products, regulating animal movement, and selective breeding for resistant traits will therefore be simpler, less expensive, and more effective to implement now than in the future. Finally, this analysis demonstrated therefore, it is recommended that these hotspots be given priority in the design and execution of BTB control methods in Ethiopia in order to lessen the disease's effects on the expanding dairy industry. The economic burden of the disease and the cost-effectiveness of these various control approaches both call for further study.

Keywords: Bovine tuberculosis, breed, prevalence, control, Ethiopia, geographic distribution

Introduction

Tuberculosis is communicable Mycobacterial disease caused by the *Mycobacterium tuberculosis* complex (MTBC) Pal (2014) ^[62]. According to recent research, *Mycobacterium* TB and *Mycobacterium bovis* have been isolated from bovine tuberculosis-infected humans and cattle, respectively (Zewed, 2014) ^[100]. Despite differences in host specificity, the members have 99.9% or more similarity at the nucleotide level and nearly identical 16S rRNA sequences (Brosch *et al.*, 2002) ^[26].

According to the OIE (2016) ^[61], the illness is a contagious chronic disease of cattle that is primarily spread between animals by inhalation, although ingestion is also frequent in cattle that graze on pastures polluted with *M. bovis*. According to Ameni *et al.* (2010) ^[12] and Ejeh *et al.* (2013) ^[34], the socioeconomic situation and low level of living in Ethiopia are more conducive to the spread of disease. Human infection occurs primarily as a result of eating undercooked meat and drinking contaminated or unpasteurized raw milk. The likelihood of transmission between cattle and humans is increased by the high frequency of TB in cattle, close contact between cattle and humans, and the custom of consuming raw milk and meat (Shitaye *et al.*, 2007) ^[108].

According to evidence, the disease reduces milk output by 10 to 18% and meat output by 15%, expenses for screening and culling infected animals, and restricts commerce (Müller *et al.*, 2013) ^[57]. Given that people and animals share the same microenvironment and living spaces, particularly in rural regions, disease in humans is currently becoming more and more relevant in poor countries. In nations where pasteurization of milk is uncommon, *M. bovis* is thought to be the cause of 10 to 15% of human tuberculosis cases (Berg *et al.*, 2015) ^[23].

Bovine TB eradication programs are in progress in several developed countries (CFSPH, 2009) ^[27]. Globally, the prevalence of the disease is estimated to be 9% based on the results of skin tests (Vorder *et al.*, 2016) ^[96] and in Ethiopia reported prevalence ranges from 3.4% (in smallholder production system) to 50% (in intensive dairy productions) and 3.5% to 5.2% in slaughterhouses (Tigre *et al.*, 2012) ^[104]. The available information is limited and

Correspondence Author:
Dr. Yeshiwas Tarekegn
HOD, Department of Animal
Health, Veterinary
Epidemiologist, Alage ATVET
College, Ziway, Ethiopia

fragmented as well as actual prevalence is at national level in the country is not well known due to inadequate disease surveillance program, socio economic impact and lack of better diagnostic facilities and fragmented articles (Asseged *et al.*, 2004) [18]. Consequently, the objective of the current systematic review is to provide.

- Current status and spatial distribution of BTB in Ethiopia based on the available reports
- To give an over view on the impact of bovine tuberculosis

Status of bovine tuberculosis in Ethiopia

According to the World Health Organization, Ethiopia is the tenth in the world, third in Africa, among the 27 MDR TB countries, and among the 22 high-burden TB countries, accounting for 81% of estimated cases (WHO, 2015) [105]. BTB in Ethiopia is most frequently detected through tuberculin skin testing, abattoir meat inspection, and rarely through bacteriological techniques. The compiled manuscripts for this review above 57 studies on PPD testing, 22 abattoir surveys paper on which all-peer reviewed were accessible online. Prevalence varies based on geographic regions, breeds, and husbandry techniques (Tschopp and Abraham, 2018) [94].

Prevalence in extensively managed production system and pastoral area

The extensive production system holds about 85% of the total livestock population of the country where as the Pastoral Production System denotes an economy that derives the bulk of its food supply from animals (milk and meat) using a variety of herding practices based on constant or partial herd mobility (oscillatory type of movement) in the low land areas of the country. Despite the presence of a huge livestock population, the actual prevalence of BTB is not known. According to Paulos (2018) [65] difficulties in sampling techniques and animal handling, combined with inadequate veterinary infrastructures are factors that hamper the process of the study. In pastoral areas in particular, the study process can be more complicated by the frequent movement of animals for water (watering points). Among the recently undertaken studies, the prevalence rate of BTB highest in Kombolcha, North shewa Zone whereas, Assella and Woldiya with the lowest. As general it ranges from 0.3 to 22.6% by in a traditionally managed extensive production system (Tschopp, 2015) [93] and at national level prevalence of 4.5% (in pastoral/agropastoral) and 4.6% (highland extensive) production system in different study area (Sibhat *et al.*, 2017) [77]. The variation may be associated with sample size, accuracy of the test and geographical difference shown below in the table.

Table 1: Prevalence reports of *B. Tuberculosis* detected by tuberculin skin test in extensively managed production system including pastoral area of Ethiopia

Area of study	No of cattle			Reference
	Tested	Positive	%	
Afar	1087	119	11	Mamo, 2013 [52]
Afar	2550	140	5.5	Sintayehu <i>et al.</i> , 2016 [78]
Assella	584	2	0.3	Tschopp, 2015 [93]
Debre-birhan	76	11	14.5	Tadele, 1998 [83]
Bako-Gazer	492	9	1.8	Tschopp, 2015 [93]
Filtu (Somali)	421	8	2	Gumi, 2011 [46]
Woldiya	620	2	0.3	Tschopp, 2015 [93]
Kombolcha	53	12	22.6	Tadele, 1998 [83]
Western Ethiopia	460	19	4.1	Laval and Ameni, 2004 [51]
Dessie	34	4	11.8	Tadele, 1998 [83]
West-wellega	353	12	3.4	Regassa, 2005
North shewa Zone	1041	169	16.2	Regassa, 2005
North shewa (Oromia)	287	27	9.4	Ayana, 2013 [19]
Meskan	624	4	0.3	Tschopp, 2015 [93]
South Ethiopia	894	36	4.0	Spiess, 2011 [81]
Total	9576	574	5.99	

Prevalence in small holder production system

Dairy animals are raised for subsistence and/or commercial milk production purposes in highland areas close to towns, where the small holder production system is predominately used. Studies on BTB prevalence in this industrial system have not been done properly. Few cross-sectional studies conducted at different geographical area of the country by using tuberculin skin tests reported the prevalence rate ranges from 3.5% in Assella (Redi, 2003) [69] to 50% in Dirediwa (Kemal, *et al.*, 2019) [49] and the study conducted

by Sibhat *et al.*, (2017) [77] at the national level shows prevalence rate of 4.6%. The prevalence in this production system increases in a chronological order which is associate with the expansion of exotic or cross breeds over the Bos tuars which agrees with Allen *et al.*, (2010) [9]. Statement Zebus are more resistant to BTB than high-producing European breeds, according to Benkirane (1998) [22], who also claimed that 48% of the variation seen in response to *M. bovis* infection is caused by host genetic diversity (Table.1).

Table 2: Based on a tuberculin test, the prevalence of *B. Tuberculosis* in small-scale dairy farms

Study Area	No of cattle			Reference
	Tested	Positive	%	
Harar	224	25	11.2	Kemal <i>et al.</i> , 2019 ^[49]
Dire diwa	58	29	50	Kemal <i>et al.</i> , 2019 ^[49]
Jigjiga	33	10	30.3	Kemal <i>et al.</i> , 2019 ^[49]
Gonder town	109	9	8.3	Shewatek, 2015 ^[74]
Mekele town	480	54	11.3	Fikre <i>et al.</i> , 2014 ^[39]
Gonder zuria	180	14	7.78	Shewatek, 2015 ^[74]
Guto Gidda District/E/wellega	295	24	8.14	Disassa <i>et al.</i> , 2016 ^[32]
Dilla town	440	19	4.3	Romha <i>et al.</i> , 2014 ^[72]
Holleta	381	25	6.4	Teshome, 1995 ^[90]
Selale	1528	18	5.1	Teshome, 1995 ^[90]
Wolayta-sodo	416	59	14.2	Regassa, 2005
Fiche	235	31	4.2	Gemta, 2000 ^[44]
Wuchale-jida	263	60	7.9	Ameni <i>et al.</i> , 2003 ^[102]
Assella	514	18	3.5	Redi, 2003 ^[69]
Total	5156	395	7.7	

Notice *Comparative intra dermal test and **Single intra dermal test

Prevalence in Intensive production system

Although the total number of cattle in this production system is small in comparison to the entire number of animals in the country, it is the primary source of milk for city residents. Better prevalence studies have been conducted in comparison to other production systems, and more frequent instances and greater prevalence rates of BTB have been noted. Taking into account the completed

tuberculin skin tests, rate of 24.3% to 65.8% (Ameni *et al.*, 2006) ^[103], and 22.9% (Ambaw, *et al.*, 2017) ^[10] and at the national level 16.6% (Sibhat *et al.*, 2017) ^[77] have been reported as we have analyzed the report on table 3 in Addis Ababa city the prevalence increases with time chronological order which is associated with expansion of dairy farms due to population overgrowth hence increments of milk and meat consumption.

Table 3: Prevalence of BTB detected by CIDT in intensive dairy farms

Area of study	No of cattle			Reference
	Tested	Positive	%	
Sululta district	858	98	11.4	Akililu <i>et al.</i> , 2014 ^[7]
Holta state farm	363	83	22.9	Ambaw <i>et al.</i> , 2017 ^[10]
Addis Ababa/C. Ethiopia/	2956	887	30	Firdessa, 2013 ^[42]
Addis Ababa	1132	386	34.1	Tsegaye <i>et al.</i> , 2010 ^[95]
Mekele	50	27	38	Abie <i>et al.</i> , 2017 ^[2]
Gonder town	28	9	17	Abie <i>et al.</i> , 2017 ^[2]
Bahir Dar town	788	10	1.27	Nuru <i>et al.</i> , 2015 ^[59]
Hawassa	22	5	11	Abie <i>et al.</i> , 2017 ^[2]
Ambo	133	37	27.8	Ameni <i>et al.</i> , 2006 ^[103]
Adama	524	58	11.1	Amin and Erkihun, 2007 ^[11]
Bako-Gazer	582	5	0.9	Tschopp <i>et al.</i> , 2015 ^[93]
Asella	281	23	8.2	Alemu, 1992
Debre-Birhan	51	3	5.9	Tadele, 1998 ^[83]
Debre-zeit state farm	114	31	27.2	Ambaw <i>et al.</i> , 2017 ^[10]
Debreziet	558	95	17.02	Meseret <i>et al.</i> , 2016 ^[55]
Adaberga state farm	243	5	2.1	Ambaw <i>et al.</i> , 2017 ^[10]
Dessie	121	89	73.6	Ameni <i>et al.</i> , 2003 ^[102]
Holleta	70	17	24.3	Ameni <i>et al.</i> , 2003 ^[102]
Kombolcha	197	96	48.7	Tadele, 1998 ^[83]
Mojo	493	338	68.6	Teshome, 1996 ^[90]
Fiche	1041	167	16	Regassa <i>et al.</i> , 2010 ^[70]
Repi	481	310	64.4	Anonymous, 1999 ^[16]
Woldiya	1029	15	1.45	Aylate <i>et al.</i> , 2013 ^[20]
Sebeta	37	4	10.8	Ameni <i>et al.</i> , 2006 ^[103]
Sellale	44	3	6.8	Ameni <i>et al.</i> , 2006 ^[103]
Ziway	205	56	27.3	Ameni <i>et al.</i> , 2003 ^[102]
Eastern Ethiopia	316	64	20.3	Kemal <i>et al.</i> , 2019 ^[49]
Total	12717	2921	23.3	

Prevalence of *Mycobacterium bovis* at Abattoir House

In Ethiopia 20-30% of cattle is slaughtered in municipal abattoirs and thus undergoes a routine meat inspection (Etter *et al.*, 2006) ^[36]. The visible tuberculosis lesions on infected cattle are observed in order to identify tuberculous lesions;

however, the standard of these procedures may differ from location to location and/or abattoir to abattoir within the nation, and studies have shown that not all *M. bovis*-infected cattle have visible tuberculous lesions at slaughter (Teklu *et al.*, 2004) ^[86]. Despite the fact that the detection of

tuberculosis lesions through abattoir inspection is currently the standard practice in Ethiopia, this may restrict the sensitivity of this abattoir detection technique. Prevalence rates range from 3.5% (Akililu *et al.*, 2014) ^[7] to 32.8%

(Biffa *et al.*, 2010) ^[24] in the investigations on abattoirs that have been conducted in different abattoirs in the country (Table 4).

Table 4: Prevalence of *B. tuberculosis* detected by abattoir meat inspection in cattle

City abattoirs	Examined	Positive	%	Reference
Sululta district	1107	39	3.5	Akililu <i>et al.</i> , 2014 ^[7]
Gambella	500	66	13.2	Alemu <i>et al.</i> 2016 ^[8]
Butajira	446	40	9.0	Biratu <i>et al.</i> , 2014 ^[25]
Addis Ababa	509	91	17.9	Biffa <i>et al.</i> , 2010 ^[24]
Nekemt	1183	70	5.9	Woyessa <i>et al.</i> , 2014 ^[99]
Adama	486	33	6.8	Terefe, 2014) ^[88]
Yabello	415	18	4.3	Biffa <i>et al.</i> , 2010 ^[24]
Hawassa	753	44	5.8	Tekle, 2016 ^[85]
Woldiya	1029	63	6.12	Tsegaye <i>et al.</i> , 2010 ^[95]
Jimma	468	35	7.5	Tesfaye, 2017 ^[89]
Melge-wendo	1265	60	4.7	Biffa <i>et al.</i> , 2010 ^[24]
Hawassa	403	88	21.8	Biffa <i>et al.</i> , 2010 ^[24]
Hossana	751	34	4.53	Teklu <i>et al.</i> , 2004 ^[36]
Kombolcha	57965	265	0.46	MoA, 2001 ^[56]
Nazareth	1125	58	5.16	Ameni and Wudie, 2003 ^[102]
Wolaita-Sodo	402	32	7.96	Regassa, 2005
Total	68807	1036	7.7	

Prevalence of *B. tuberculosis* based on breed of animals at the national level

Frequency of BTB in Ethiopian cattle. According to the results section below, prevalence of 21.6% for Holstein-Friesian cattle and 9.9% for crosses were significantly higher than the prevalence of 4.1% for zebu cattle. This supports the idea that genetic factors may influence a cattle's susceptibility or resistance to BTB (Finaly *et al.*, 2012) ^[40]. Studies done in Ethiopia in this area showed that Holstein-Friesian cattle had higher prevalence and severity of TB lesions than zebu cattle and their crosses kept together on communal pasture in the country's central highlands (Ameni *et al.*, 2007 ^[11] as shown in table 5). Similar results were

found in an experimental study comparing the susceptibility of Holstein-Friesian calves to Boran (zebu) calves exposed to low doses of *M. bovis* in South Africa. The study found that while none of the Boran calves developed BTB suggestive lesions, 50% of the Holstein-Friesian calves developed typical lesions. The results showed that zebu breeds, as opposed to the Holstein-Friesian breed, are more likely to be resistant to BTB infection. Zebus may be more resistant to BTB than high-yielding European breeds, according to Benkirane (1998) ^[22]. The entirety of this study supports the claim made by Allen *et al.* (2010) ^[9] that 48% of the variance observed in the host's response to *M. bovis* infection is caused by genetic variation.

Table 5: Prevalence of *B. Tuberculosis* based on breed of animals at the national level

Study area	Breed						Reference
	Zebu	Cross	Holstein	Nuer	Felata	Horo	
Ethiopia	4.1	9.9	21.6	-	-	-	Sibhat <i>et al.</i> , 2017 ^[77]
Sululta district	1.8	12.8	-				Akililu <i>et al.</i> , 2014 ^[7]
Gambella	-	-	-	12.9	57.1	10.3	Alemu, 2015
Nekemit	5.7	6.7	-	-	-	-	Woyessa <i>et al.</i> , 2014 ^[99]
Bahirdar town	0.2	2.38	-	-	-	-	Nuru <i>et al.</i> , 2015 ^[59]
Dilla town	1.7	3.8	9.6	-	-	-	Romha <i>et al.</i> , 2014 ^[72]
Gonder town	0	11.7	-	-	-	-	Shewatek, 2015 ^[74]
E/wellega	1.75	9.6	-	-	-	-	Disassa <i>et al.</i> , 2016 ^[32]
Hawassa	4.4	17.9	-	-	-	-	Tekle, 2016 ^[85]
Mekele	2.7	15.8	14.8				Fikire <i>et al.</i> , 2014 ^[39]
Jimma	5.02	43.3	-	-	-	-	Tesfaye, 2017 ^[89]

Prevalence *B. tuberculosis* based on Geographical Sites at the national level

In general, the prevalence of BTB decreased as one moved from the country's center to its periphery in all directions, as well as from easily accessible regional towns to more distant regions where traditional intensive livestock agriculture predominates. The generalization is not unqualified because variations may be seen over close distances or even in the same region, as recorded by various researchers at various times. The prevalence of BTB appears to be highest in the country's center, with Addis Abeba and the adjacent sub-

urban areas experiencing the highest intensity. This may be the result of the relatively long history of intensive dairy farming in central Ethiopia using exotic European breeds that were introduced during the previous imperial regime with the aim of supplying dairy products to the expanding human population in Addis Abeba and its surroundings (Ahmed *et al.*, 2004) ^[5]. These regions have a significant concentration of dairy cattle, and they provide the rest of the nation with enhanced dairy animals as well as BTB for the developing ones (Firdessa *et al.*, 2012) ^[41]. It also appears that most studies have been performed in areas adjacent to

existing roads, evidently for logistical reasons. Most rural areas further away from road access have not been investigated at all that includes most cross-border areas where there is a high informal animal movement for instance between Ethiopia and Sudan, Kenya or Somalia (Tschopp and Abraham, 2018) [96].


Geographical distribution of *B. tuberculosis* based on administrative state

The frequency of BTB does not significantly differ amongst the various administrative regions of the nation. This is because none of the administrative states have implemented

any workable control measures, therefore the epidemiology of BTB is unaffected by animal health operations in any of them. As there are no active regulations limiting animal movement due to biosecurity, the results of this research indicate that intensifying the dairy industry could increase prevalence in any of the locations. This has a negative impact on the dairy industry because traffic in dairy cows was found to be unidirectional in Ethiopia and was often from the region's central regions, where BTB is more common, to the zonal towns and regional cities (Firdessa *et al.*, 2012) [41]. The rest regional states have no valid general prevalence which recalls for further studies in these regions.

Table 6: Prevalence reports of *Bovine tuberculosis* administrative state

Administrative state	Prevalence (95% confidence interval)	Reference
Addis ababa	10.6 (6.7-16.4)	Sibhat <i>et al.</i> , 2017 [77]
Tigray	8.8 (5.8-13.3)	
Afar	6.7 (3.4-12.7)	
Oromia	6.6 (5.5-8.1)	
SNNP	4 (2.9-5.6)	
Amhara	3.6 (2.2-5.8)	



Source: Sibhat *et al.*, 2017 [77]

Fig 2: Geographical Distribution of *Bovine tuberculosis* in Ethiopia

Economic and zoonotic importance of *Bovine Tuberculosis* in Ethiopia

Because of differences in epidemiological conditions, livestock systems, natural reservoirs, time horizons, and the absence of generally accepted analytical frameworks, the global economic assessment of loss associated with BTB and cost-benefits analysis from its control were multifaceted. Data obtained from different countries provide variable results. The illness significantly affects the amount of milk and meat an affected cow produces, and it also has

an impact on animal reproductive and pulling strength in conventional farming systems (Zinsstag *et al.*, 2006) [101]. Additionally, the economic and financial effects of tuberculosis on society's healthcare expenses. Since 75% of TB patients fall within the 15–54 age range, which is an economically active period, the disease presents a barrier to socioeconomic progress. Globally for Control, surveillance and monitoring costs annual agricultural loss USD 3 billion and BTB eradication in the US did cost 538 million USD between 1917 and 1992 (Nelson 1999) [58] also costing

currently in the UK 100 million pound per year (Matthew *et al.*, 2006) ^[54].

The disease's economic repercussions in Ethiopia have not yet been thoroughly explored or reported. There haven't been many abattoir meat inspection surveillances that reveal the percentage of complete or partial organ and carcass condemnations. Gezahegne (1991) ^[45] provided evidence that the condemned carcasses and organs from 1.2 million slaughtered cattle in eight export abattoirs resulted in an estimated cost of more than 300,000,000,000 US dollars or 600,000,000 ETB. Recently, Tschopp *et al.* (2012) ^[106] calculated that between 2005 and 2011, the economic burden of BTB ranged from 500,000 to 4.9 million US dollars in urban livestock production systems and from 75.2 million and 385 million US dollars in vast rural animal production. These numbers showed losses of less than 1% of the livestock's net present value in the rural and 3.9–6.2% in urban livestock production systems per year. (Sibhat *et al.*, 2017) ^[77].

For non-economic reasons, such as concerns over the spread of BTB through the trade of dairy cattle from a system with a high prevalence of the disease to one with a low prevalence of sedentary lifestyles, as well as for public health, Tschopp *et al.* (2012) ^[106] emphasized the urgent need for control of the disease in the urban production system in Ethiopia. Worldwide, zoonotic tuberculosis in cattle is a problem for public health. High levels of HIV and poverty, particularly in Sub-Saharan nations, are major factors in the developing world. According to Cezar *et al.* (2016) ^[28], consumption of raw milk or undercooked milk products is the primary cause of non-pulmonary tuberculosis. Risk factors include social-cultural practices, such as the custom of pastoral tribes to consume raw blood, raw milk, raw or undercooked meat, and meat products (Amenie and Wudie, 2003) ^[102]. According to estimates, 90% of the milk drank in Africa is either consumed raw or fermented, increasing the risk of disease transmission (Ibrahim, 2012). According to Tigre *et al.* (2011) ^[92], tuberculosis (TB) infects 9.4 million people worldwide and results in 1.4 million annual fatalities. Asia and Africa combined for 55% and 30% of all TB cases worldwide, respectively. Eastern Mediterranean (7%), European (4%), and American (3%), regions saw lower percentages of instances. Following malaria and births as the third and fourth top causes of hospital admissions, it is also the second biggest cause of death. (Regassa *et al.* 2010) ^[70].

Bovine tuberculosis Control options

Bovine tuberculosis is listed under the OIE terrestrial animal health code, and control should be aimed at reducing prevalence in animals in order to prevent transmission to humans. In Ethiopia, control measures cannot be adopted in practice due to a variety of factors, including: lack of knowledge on the actual prevalence of the disease, the existing technical and financial limitations, a lack of veterinary infrastructures, cultural and/or traditional beliefs, and geographical barriers.

Control in the Cattle Populations

Due of the numerous vulnerable species, diverse pathophysiology, and poor efficacy of currently used methods for wild animals, *M. bovis* is challenging to control. Applying the following management strategies needs an awareness of the epidemiology of infection within

the ecological system, which might involve both domestic and wild animal species (Cousins, 2001) ^[30]. The only method that guarantees eliminating up to testing every three months to eliminate individuals in the herd that can spread infection is the test and slaughter method. It is beneficial to take routine hygiene precautions such cleaning and disinfecting polluted areas, food, and water troughs. According to Aiello *et al.* (1998) ^[6], cattle under poor management were more likely to contract tuberculosis than cattle under competent management. In developing nations, which cannot afford a test and slaughter control program, particularly in nations with a wild life reservoir of *M. bovis* infection, vaccination against tuberculosis is expected to become a major disease control method. Significant advancements have been made in the creation and testing of TB vaccines for cattle and a variety of wild animals over the past ten years (Edelsten, 1999) ^[33].

Control in the human populations

In general, the BCG vaccine and the use of chemotherapy can reduce the risk of both humans and animals contracting tuberculosis. The common anti-tuberculosis medications, including ionized, rifampicin, pyrazinamide, thiacetazone, and ethambutol, are used to treat patients and stop the disease from spreading. The length of the treatment program might be either brief (two months) or standard (six to eight months). Only 76% of new patients in the 2002 cohort experienced treatment success, which is significantly less than the maximum recorded success rate of 80% for Ethiopia (WHO, 2005) ^[107]. Additionally, health education is being used as one of the key methods for controlling through sanitization and raising community understanding of the epidemiological characteristics of the disease. Other efficient steps are also being taken to enable improved access across the nation.

Conclusion and recommendation

In comparison to the prevalence seen in the established dairy belt in the central areas of Ethiopia, particularly in and around Addis Ababa City, this review revealed comparatively low average prevalence in the emerging dairy districts. According to the findings of skin tests, the estimated prevalence of *B. tuberculosis* in Ethiopia was determined to be 5.8%, which is lower than the estimated global prevalence (9%) of the disease. Due to the current lower number of infected animals in a more constrained geographic area, implementing a control program in these cities could be simpler, cheaper, and more effective to do so now than in the future. Test and slaughter programs at this stage are not economically feasible nationwide in Ethiopia. There is an urgent need to investigate alternative cheaper options for BTB control such as milk pasteurization meat inspection, awareness regulation in animal movement control and selective breeding program for resistant traits are optional. More research is warranted regarding the economic impact of the disease (productivity losses in animals, public health impacts, social impacts, household micro-economics, and market losses) and cost-efficiency of these different control options. On the basis of these, review the following recommendations are forwarded

- To know accurate prevalence, distribution and better insight into the transmission scenario of the disease further study on larger sample size, molecular level and nationwide epidemiological survey should be

conducted

- In the majority of Ethiopia, animals are housed close to homes and kept in extremely unhygienic conditions, also consume badly processed meat and milk products, so raising awareness of this issue is vital.
- Following applying for insurance, dairy producers may be persuaded to kill their sick cattle after testing for BTB and other economically significant infectious illnesses.
- The diagnosis of tuberculosis lesions requires routine tuberculin skin testing and standard abattoir meat inspection processes, and the outcome can be improved by using Ziehl-Neelsen staining concurrently.

References

1. Abel T. Survey of abattoir inspection data in Dire-Dawa abattoir. Annual Report, Dire-Dawa, Eastern Ethiopia; c1989.
2. Abie G, Andrew B, Birhanu T, Alemseged A, Sintayehu G, Biniam T, Ameni G. Prevalence of *B. tuberculosis* and its associated risk factors in Ethiopia. *Journal of Preventive Veterinary Medicine*; c2017. <https://doi.org/10.1016/>.
3. Acha P, Szytles B. Zoonotic tuberculosis, zoonosis and communicable diseases to man and animals, 8th ed. Washington D.C. 20037. USA; c2001.
4. Admasu P, Berihun W, Niguse A. Prevalence of bovine tuberculosis in dairy cattle of Yeki District, Southern Ethiopia. *Afr J Basic Appl Sci*. 2014;6(5):135-40.
5. Ahmed M, Ethui S, Yemesrach A. Dairy development in Ethiopia (EPTD). Discussion paper No. 123 (International Food Research Institute) Washington; c2004.
6. Aiello SE, Mays A. The Merck Veterinary Manual. Whitehouse Station, NJ: Merck & Co. Inc. 1998;8:2187-97.
7. Akililu B, Ameni G, Teshale S, Fanta D, Akafte T, Ketema T. Epidemiology and public health significance of *Bovine tuberculosis* in and around sululta district central Ethiopia. *African journal of microbiology research*. 2014;8(24):2352-2358.
8. Alemu T, Mamo G, Ameni G, Pal M. Molecular epidemiology of *B. tuberculosis* in cattle and its public health implications in gambella region. *Molecular Microbiol. Res*. 2016;6(1):1-15.
9. Allen AR, Minozzi G, Glass EJ, Skuce RA, McDowell SW, Woolliams JA, *et al.* Bovine tuberculosis: the genetic basis of host susceptibility. *Proceedings of the Royal Society B: Biological Sciences*. 2010 Sep 22;277(1695):2737-45.
10. Ambaw M, Benti D, Gobena A. *Bovine Tuberculosis* Prevalence, Potential Risk Factors and Its Public Health Implication in Selected State Dairy Farms, Central Ethiopia. *World Vet J*. 2017;7(1):21-29.
11. Ameni G, Erkihun A. Bovine tuberculosis on small-scale dairy farms in Adama Town, central Ethiopia, and farmer awareness of the disease. *Revue Scientifique et Technique-Office International des Epizooties*. 2007 Dec 1;26(3):711-20.
12. Ameni G, Desta F, Firdessa R. Molecular typing of *Mycobacterium bovis* isolated from tuberculosis lesions of cattle in north eastern Ethiopia. *Veterinary Record*. 2010 Jul;167(4):138-41.
13. Ameni G, Vordermeier M, Firdessa R, Aseffa A, Hewinson G, Gordon SV, *et al.* *Mycobacterium tuberculosis* infection in grazing cattle in central Ethiopia. *The Veterinary Journal*. 2011 Jun 1;188(3):359-61.
14. Amit K, Ruchi T, Sandip C, Neha M, Kuldeep D, Shoorgh B. In-sight into Bovine tuberculosis, various approaches for Diagnosis, control and its public health concerns. *Asian Journal of Animal and veterinary Advances*. 2014;9(6):324-344.
15. Anonymous. Zoonotic tuberculosis: a memorandum from World Health Organization with participation of Food and Agricultural Organization (FAO). *Bulletin of the WHO*. 1994;72:851-857.
16. Anonymous. Dairy development enterprise. Annual report Addis Ababa, Ethiopia; c1999.
17. Ashford D, Whitney E, Raghunath P, Cosivi O. Epidemiology of *M.bovis* that infect humans and animals, Scientific and Technical Review, OIE. 2001;20:325-337.
18. Asseged Z, W/dsenbet E, Yimer E. Evaluation of abattoir inspection for the diagnosis of *M. bovis* Addis Ababa. *Trop. Anim. Health Prod*. 2004;36:537-546.
19. Ayana T, Tafesse K, Mamo G, Tessema T, Ameni G. Isolation and molecular characterization of non-tuberculosis mycobacteria from skin positive reactors and pathological lesions of cattle at Bako, Western Ethiopia. *AJMR*. 2013;20:2190-2197.
20. Aylate A, Shah H, Aleme T, Gizaw T. Bovine tuberculosis: prevalence and diagnostic efficacy of routine meat inspection procedure in Woldiya municipality abattoir North Wollo zone, Ethiopia. *Trop. Anim. Health Prod*. 2013;45:855-864.
21. Bekele M, Indris B. Evaluation of Routine Meat Inspection Procedure to Detect *Bovine Tuberculosis*, Suggestive Lesions in Jimma Municipal Abattoir, South West Ethiopia. *Global Veterinaria*. 2011;6(2):172-179.
22. Benkirane A. Bovine Tuberculosis in Africa. *World Animal Review*; 1998, 90-1998/<http://www.fao.org/docrep/W8600T/w8600t09.htm> Google Scholar.
23. Berg S, Schelling E, Hailu E, Firdessa R, Gumi B, Erenso G, *et al.* Investigation of the high rates of extrapulmonary tuberculosis in Ethiopia reveals no single driving factor and minimal evidence for zoonotic transmission of *Mycobacterium bovis* infection. *BMC infectious diseases*. 2015 Dec;15(1):112.
24. Biffa D, Skjerve E, Oloya J, Bogale A, Abebe F, Dahle U, *et al.* Molecular characterization of *Mycobacterium bovis* isolates from Ethiopian cattle. *BMC Veterinary Research*. 2010 Dec;6(1):1-1.
25. Biratu N, Gebremedhn G, Tadesse B, Habitamu T, Gebrehiwot T, Belayneh G. epidemiology of Bovine tuberculosis in Butajira, Southern Ethiopia: cross-sectional abattoir based study. *African journal of microbiology research*. 2014;8(33):3121-3117.
26. Brother S, White I, Coffey S, Downs A, Mitchell R, Good J. Evidence of genetic resistance of cattle to infection with *M. bovis*, *J. Dairy Sci*. 2010 Mar 1;93(3):1234-42.
27. Center of Food Security and Public Health and Instituion of cooperation in animal biology. *Bovine Tuberculosis* Iowa state university; c2009. Available at H:\CFSPH\CFSPH materials /CFSPH, Disease Otemplates/ Tech Fs /Fact -Sheet -1090-Bar-2012.

28. Cezar RD, Lucena-Silva N, Borges JM, Santana VL, Junior JW. Detection of *Mycobacterium bovis* in artisanal cheese in the state of Pernambuco, Brazil. *International Journal of Mycobacteriology*. 2016 Sep 1;5(3):269-72.
29. Cosivi O, Grange JM, Daborn CJ, Raviglione MC, Fujikura T, Cousins D, *et al.* Zoonotic tuberculosis due to *Mycobacterium bovis* in developing countries. *Emerging infectious diseases*. 1998 Jan;4(1):59.
30. Cousins DV. *Mycobacterium bovis* infection and control in domestic livestock. *Revue scientifique et technique (International Office of Epizootics)*. 2001 Apr 1;20(1):71-85.
31. Demelash B, Inangolet F, Oloya J, Asseged B, Badaso M, Yilkal A, *et al.* Prevalence of bovine tuberculosis in Ethiopian slaughter cattle based on post-mortem examination. *Tropical animal health and production*. 2009 Jun;41:755-65.
32. Disassa H, Mezene W, Tadesse B, Sultan A, Fikadu B, Ketema T. A Cross-sectional Study on *Bovine Tuberculosis* in Small Holder dairy farms of Guto Gidda District, East Wollega Zone, Western Ethiopia. *Nature and Science*. 2016;14(3). <http://www.sciencepub.net/nature>.
33. Edelsten R. Tuberculosis in cattle in Africa, control measures and implication for human health. *Proceeding of International Conference at the International livestock research Institute (ILRI)*. Addis Ababa, Ethiopia; c1999.
34. Ejeh EF, Markus IF, Ejeh AS, Musa JA, Lawan FA, Ameh JA, *et al.* Seasonal prevalence of bovine tuberculous lesions in cattle slaughtered in Yola abattoirs. *Bangladesh Journal of Veterinary Medicine*. 2013 Dec 1;11(2):113-20.
35. Elias K, Hussein B, Asseged T, Wondwossen M, Gebeyehu. Status of Bovine tuberculosis in Addis Ababa dairy farms *Rev. Sci. Tech*. 2008;27(3):915-923.
36. Etter E, Ameni G, Roger F. Tuberculosis: risk assessment in Ethiopia: safety of meat from cattle slaughtered in abattoirs. *Proceedings of the 11th International Symposium on Veterinary Epidemiology and Economics*, Cairns, Australia; c2006.
37. Fetene T, Kebede N. Bovine tuberculosis of cattle in three districts of north western Ethiopia. *Tropical Animal Health and Production*. 2009;41:273-277.
38. Figueroa-Munoz J, Ramon-Pardo P. Tuberculosis control in vulnerable groups. *Bulletin of the World Health Organization*. 2008;86:657-736.
39. Fikre Z, Romha G, Berhe G, Mamo G, Sisay T, Ameni G. Prevalence of *Bovine tuberculosis* and assessment of Cattle owners' awareness on its public health implication in and around Mekelle, Northern Ethiopia. *Journal of veterinary medicine and Animal health*; c2014. DOI: 10.5897/JVMAH2014:288 Article no 48961c944606.
40. Finlay E, Berry D, Wickham B, Gormley E, Bradley G. A genome wide association scan of Bovine tuberculosis susceptibility in Holstein-Friesian dairy cattle, *PLoS One*. 2012;7(2):e30545, 10.1371/journal.pone.0030545 CrossRefGoogle Scholar.
41. Firdessa R, Tschopp R, Wubete A, Sombo M, Hailu E, Kiros T, *et al.* High prevalence of *B. tuberculosis* in dairy cattle in central Ethiopia: implications for the dairy industry and public health. *PLoS One*. 2012;7(12):e52851, 10.1371/journal.pone.0052851.
42. Firdessa R, Berg S, Hailu E, Schelling E, Gumi B, Erenso G, Gadisa E, Kiros T, Habtamu M, Hussein J, Zinsstag J. Mycobacterial lineages causing pulmonary and extrapulmonary tuberculosis, Ethiopia. *Emerging infectious diseases*. 2013 Mar;19(3):460.
43. GebreMariam S, Amare S, Baker D, Solomon A, Davies R. Study of the Ethiopian live cattle and beef value chain. *ILRI Discussion Paper 23*. Nairobi: International Livestock Research Institute; c2013.
44. Gemta M. A cross-sectional study on *Bovine tuberculosis* in small holder dairy farms and implication in man in, Fitch town, North Shewa. *DVM Thesis*. Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia; c2000.
45. Gezahegne L. Economical aspect of condemned organs and parts due to cysticercosis, hydatidosis, fasciolosis and tuberculosis. *Analysis report MoA*; c1991.
46. Gumi B, Schelling E, Firdessa R, Aseffa A, Tschopp R. Prevalence of *Bovine tuberculosis* in pastoral cattle herds in the Oromia region, Southern Ethiopia. *Tropical Animal Health and Production*. 2011;43(6):1081-1087.
47. Higgins J, Camp P, Pate M, Robbe S. Identification of *Mycobacterium* spp. of veterinary importance using *rpoB* gene sequencing. *BMC Vet Res*. 2011;7:77.
48. Ibrahim S, Cadmus S, Umoh J, Ajogi I, Farouk U, Abubakar U. Tuberculosis in humans and cattle in Jigawa state, Nigeria: Risk factors analysis. *Veterinary Medicine International*; c2012. DOI: 10.1155/2012/865924
49. Kemal J, Sibhat B, Abraham A, Terefe Y, Tulu KT, Welay K, *et al.* Bovine tuberculosis in eastern Ethiopia: prevalence, risk factors and its public health importance. *BMC Infectious Diseases*. 2019 Dec;19(1):1-9. <https://doi.org/10.1186/s12879-018-3628-1>.
50. Kiros. Epidemiology and zoonotic importance of *Bovine tuberculosis* in selected sites of Eastern Shewa Ethiopia. *MSc. Thesis*. Faculty of Veterinary Medicine, Addis Ababa, University and Freie Universitat, Berlin, Germany; c1998.
51. Laval G, Ameni G. Prevalence of *B. tuberculosis* in zebu cattle under traditional animal husbandry in Boji district of Ethiopia. *Revue Méd. Vét*. 2004;155(10):494-499.
52. Mamo G, Abebe F, Worku Y, Hussein N, Legesse M, Tilahun G, *et al.* Bovine tuberculosis and its associated risk factors in pastoral and agro-pastoral cattle herds of Afar Region, Northeast Ethiopia. *J Vet Med Anim Health*. 2013 Jun;5(6):171-9.
53. Marie-France H, Maria L, Claude S. Classification of worldwide *Bovine tuberculosis* risk factors in cattle: a stratified approach. *Vet Res*. 2009;40:50.
54. Mathews F, Macdonald DW, Taylor GM, Gelling M, Norman RA, Honess PE, *et al.* Bovine tuberculosis (*Mycobacterium bovis*) in British farmland wildlife: the importance to agriculture. *Proceedings of the Royal Society B: Biological Sciences*. 2006 Feb 7;273(1584):357-65.
55. Meseret B, Gezahegn M, Samuel M, Gobena A, Gashaw B, Etsegenet T. Epidemiology of *Bovine Tuberculosis* and Its Public Health Significance in Debre- Zeit Intensive Dairy Farms, Ethiopia. *Biomedicine and Nursing*; c2015.

- <http://www.nbmedicine.org>.
56. MoA. Pastoral and agro pastoral areas extensionsystem. Agricultural Extension Department, Ministry of Agriculture, Addis Ababa, Ethiopia; c2001.
 57. Müller B, Dürr S, Alonso S, Hattendorf J, Laisse CJ, Parsons SD, *et al.* Zoonotic *Mycobacterium bovis*-induced tuberculosis in humans. *Emerging infectious diseases*. 2013 Jun;19(6):899-908.
 58. Nelson AM. The cost of disease eradication: smallpox and bovine tuberculosis. *Annals of the New York Academy of Sciences*. 1999 Dec;894(1):83-91.
 59. Nuru A, Mamo G, Teshome L, Zewdie A, Medhin G, Pieper R, *et al.* Bovine tuberculosis and its risk factors among dairy cattle herds in and around Bahir Dar City, Northwest Ethiopia. *Ethiopian Veterinary Journal*. 2015 Dec 3;19(2):27-40.
 60. Nwanta J, Joseph I, Ezema W, Sunday, Umenonig C. tuberculosis: A review of epidemiology, clinical presentation, prevention and control. *Journal of public Health and Epidemiology* 2010 Sep;2(6):118-24.
 61. OIE. OIE Manual of Diagnostic Tests and Vaccines for Terrestrial Animals Office Internationale des Epizooties (OIE), Paris, France Google Scholar; c2016.
 62. Pal M, Zenebe N, Rahman M. Growing significance of *Mycobacterium bovis* in human health. *Microb H*. 2014;3:29-34.
 63. Palmer M, Thacker T, Corner L. *M. bovis*: A model pathogen at the Interface of Livestock and Humans. *Veterinary Medicine International*. 2012;1:17.
 64. Panarella ML, Bimes RS. A naturally occurring outbreak of tuberculosis in a group of imported cynomolgus monkeys (*Macaca fascicularis*). *Journal of the American Association for Laboratory Animal Science*. 2010 Mar 15;49(2):221-5.
 65. Paulos C. Review on Status of *B.tuberculosis* in Ethiopia, DVM seminar paper Jimma University College of Agriculture and Veterinary Medicine, Jimma, Ethiopia; c2018.
 66. Quinn P, Carter M, Markey B, Carter G. *Mycobacterium* species. *Clinical veterinary microbiology*. London Philadelphia, 1999, 157-170.
 67. Radostits O, Gay C, Hinchcliff K, Constable P. A textbook of the diseases of cattle, horses, sheep, pigs and goats 10th ed. Saunders, Toronto, Canada, 2006, 1007-1015.
 68. Radostits O, Gay C, Hincheliff K, Constabel P. *Vet. Medicine*. A text book of the disease of cattle, shoat, pig, and horses. 10th ed., Elsevier, London; c2007. p. 1007-1040.
 69. Redi N. Prevalence of *Bovine tuberculosis* and zoonotic implication in Asela Town, South East Ethiopia. *DVM Thesis*. Faculty of Veterinary Medicine, Addis Ababa University, Debre-Zeit, Ethiopia; c2003.
 70. Regassa A, Tassew K, Amenu B, Megersa F, Abunna B Mekibib, Amneni G. A cross-sectional study on Bovine tuberculosis in Hawassa town and its surroundings, Ethiopia. *Tropical Animal Health and Production*. 2010;42:915-920.
 71. Ritacco V, Torres P, Sequeira M, Reniero A, De Kantor I. *Bovine Tuberculosis* in Latin America and Caribbean: Infections in Animals and Humans. 2nd edth. Wiley Publishers, Hoboken, N.J., USA; c2008. p. 149-160.
 72. Romha G, Gebreegziabher G, Ameni G. Assessment of bovine tuberculosis and its risk factors in cattle and humans, at and around Dilla town, southern Ethiopia. *Animal Vet Sci*. 2014;2(4):94-100.
 73. Senedu B, Solomon A, Gunnar A. Federal Ministry of Health Ethiopia Tuberculosis, Leprosy and TB/HIV Prevention and Control Program Manual FMOH, Addis Ababa. *Journal of Tuberculosis Research*. 2008;4(1).
 74. Shewatatek F. Bovine Tuberculosis: Prevalence, Economic Implications And Diagnostic Test Comparison In Smallholder Dairy Farms In Selected Districts Of North Gondar Zone, Amhara Region, Ethiopia. *Msc Thesis of Veterinary Epidemiology and Economics*; c2015. www.academia.edu.
 75. Shitaye JE, Getahun B, Alemayehu T, Skoric M, Trembl F, Fictum P, *et al.* A prevalence study of bovine tuberculosis by using abattoir meat inspection and tuberculin skin testing data, histopathological and IS6110 PCR examination of tissues with tuberculous lesions in cattle in Ethiopia. *VETERINARNI MEDICINA-PRAHA*. 2006 Nov 1;51(11):512.
 76. Sennuga SO, Lai-Solarin WI, Adeoye WA, Alabuja FO. Extension's role in improving livestock production: information needs, institutions and opportunities. *Int. J Agric. Nutr.* 2022;4(2):43-51. DOI: 10.33545/26646064.2022.v4.i2a.80
 77. Sibhat B, Asmare K, Demissie K, Ayelet G, Mamo G, Ameni G. Bovine tuberculosis in Ethiopia: a systematic review and meta-analysis. *Preventive veterinary medicine*. 2017 Nov 1;147:149-57.
 78. Sintayehu W, Ignas M, Herbert H, Fitsum A, Daniel A, Zelalem E, *et al.* Risk Factors for *Bovine Tuberculosis* in Cattle in Ethiopia. *Plos one*; c2016. doi:10.1371/journal.pone.0159083.
 79. Sisay S, Mengistu B, Erku W, Woldeyohannes D. Directly Observed Treatment Short-course (DOTS) for tuberculosis control program in Gambella Regional State, Ethiopia: ten years' experience. *BMC research notes*. 2014 Dec;7(1):44.
 80. Smith NH, Gordon SV, de la Rua-Domenech R, Clifton-Hadley RS, Hewinson RG. Bottlenecks and broomsticks: the molecular evolution of *Mycobacterium bovis*. *Nature Reviews Microbiology*. 2006 Sep 1;4(9):670-81.
 81. Spiess M. *Mycobacteria* and zoonoses among pastoralists and their livestock in South-East Ethiopia, PhD thesis 2011, Swiss Tropical & Public Health Institute; c2011.
 82. Srivastava K, Chauhan DS, Gupta P, Singh HB, Sharma VD, Yadav V, *et al.* Isolation of *Mycobacterium bovis* & *M. tuberculosis* from cattle of some farms in north India-possible relevance in human health. *Indian Journal of Medical Research*. 2008 Jul 1;128(1):26-31.
 83. Tadele A. Evaluation of diagnostic tests, prevalence and zoonotic importance of BTB in Ethiopia. *DVM Thesis*. FVM, Addis Ababa University, Debre-Zeit, Ethiopia; c1998.
 84. Tamiru F, Hailemariam M, Terfa W. Preliminary study on prevalence of bovine tuberculosis in cattle owned by tuberculosis positive and negative farmers and assessment of zoonotic awareness in Ambo and Toke Kutaye districts, Ethiopia. *Journal of Veterinary Medicine and Animal Health*. 2013;5(10):288-95.
 85. Tekle Y. Isolation, and Molecular Characterization of *M. bovis* Isolated from Cattle Slaughtered at Hawassa University and Municipal Abattoirs, S. Ethiopia.

- MVSc. Thesis; c2016.
86. Teklu A, Asseged B, Yimer E, Gebeyehu M, Woldesenbet Z. Tuberculous lesions not detected by routine abattoir inspection: the experience of the Hossana municipal abattoir, southern Ethiopia. *Revue Scientifique et Technique-Office International Des Epizooties*. 2004 Dec 1;23(3):957-64.
 87. Tenguria RK, Khan FN, Quereshi S, Pandey A. Epidemiological study of zoonotic tuberculosis complex (Ztbc). *World Journal of Science and Technology*. 2011;1(3):31-56.
 88. Terefe D. Gross pathological lesions of *B. tuberculosis* and efficiency of meat inspection procedure to detect infected cattle in Adama municipal abattoir. *JVMAH*. 2014 Feb 28;6(2):48-53.
 89. Tesfaye W. study on epidemiology of *Bovine tuberculosis* and its public health Significance in jimma town and its surroundings, S/W. Ethiopia; c2017. MVSc. Thesis.etd.aau.et
 90. Teshome Y. Status of *Bovine tuberculosis* in Ethiopia: *Mycobacterium bovis* Infection in Animals and Humans. Iowa State University Ames, Iowa, USA, 1995, 273–279.
 91. Thoen CO, Kaplan B, Thoen TC, Gilsdorf MJ, Shere JA. Zoonotic tuberculosis: A comprehensive one health approach. *Medicina (Buenos Aires)*. 2016 Jun;76(3):159-65.
 92. Tigre W, Alemayehu G, Abetu T, Deressa B. Preliminary study on public health implication of bovine tuberculosis in Jimma Town, South Western Ethiopia. *Global Veterinaria*. 2011;6(4):369-73.
 93. Tschopp R, Bekele S, Moti T, Young D, Aseffa A. Brucellosis and bovine tuberculosis prevalence in livestock from pastoralist communities adjacent to Awash National Park, Ethiopia. *Preventive veterinary medicine*. 2015 Jun 15;120(2):187-94.
 94. Tschopp R, Abraham A. *Bovine tuberculosis* and other *Mycobacterium* in animals in Ethiopia: a systematic review. *Jacobs Epidemiol Prevent*. 2018;4(1):026.
 95. Tsegaye W, Aseffa A, Mache A, Mengistu Y, Stefan B, Ameni G. Conventional and Molecular Epidemiology of Bovine Tuberculosis in Dairy Farms in Addis Ababa City, the Capital of Ethiopia. *The Journal of Applied Research in Veterinary Medicine*. 2010 Jan 1;8(2):143-145.
 96. Vordermeier M, Ameni G, Berg S, Bishop R, Robertson BD, Aseffa A, Hewinson RG, Young DB. The influence of cattle breed on susceptibility to bovine tuberculosis in Ethiopia. *Comparative immunology, microbiology and infectious diseases*. 2012 May 1;35(3):227-32.
 97. Vordermeier HM, Jones GJ, Buddle BM, Hewinson RG, Villarreal-Ramos B. Bovine tuberculosis in cattle: vaccines, DIVA tests, and host biomarker discovery. *Annual review of animal biosciences*. 2016 Feb 15;4:87-109.
 98. World Organization for Animal Health (OIE): *B. Tuberculosis OIE Terrestrial Manual*; c2015. http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.04.06_B-tb.
 99. Woyessa M, Jibril Y, Ameni G, Duguma R. Molecular epidemiology of *Mycobacterium tuberculosis* complex at Nekemte municipality abattoir, Western Ethiopia. *Science, Technology and Arts Research Journal*. 2014 Aug 25;3(1):167-73.
 100. Zeweld SW. Cultural and molecular detection of zoonotic tuberculosis and its public health impacts in selected districts of Tigray region, Ethiopia. *Sokoto Journal of Veterinary Sciences*. 2014 Apr 23;12(1):1-2.
 101. Zinsstag J, Schelling E, Roth F, Kazwala R, Thoen CO, Steele JH. Economics of bovine tuberculosis. *Mycobacterium bovis* infection in animals and humans. 2006 Apr 3;2:68-83.
 102. Ameni G, Amenu K, Tibbo M. Bovine tuberculosis: prevalence and risk factor assessment in cattle and cattle owners in Wuchale-Jida district, Central Ethiopia. *Journal of applied research in veterinary medicine*. 2003.
 103. Ameni G, Aseffa A, Engers H, Young D, Hewinson G, Vordermeier M. Cattle husbandry in Ethiopia is a predominant factor affecting the pathology of bovine tuberculosis and gamma interferon responses to mycobacterial antigens. *Clinical and Vaccine Immunology*. 2006 Sep;13(9):1030-6.
 104. Tigre RC, Silva NH, Santos MG, Honda NK, Falcao EP, Pereira EC. Allelopathic and bioherbicidal potential of *Cladonia verticillaris* on the germination and growth of *Lactuca sativa*. *Ecotoxicology and environmental safety*. 2012 Oct 1;84:125-32.
 105. World Health Organization. Trends in maternal mortality: 1990-2015: estimates from WHO, UNICEF, UNFPA, World Bank group and the United Nations population division. World Health Organization; 2015.
 106. Tschopp MA, Solanki KN, Gao F, Sun X, Khaleel MA, Horstemeyer MF. Probing grain boundary sink strength at the nanoscale: Energetics and length scales of vacancy and interstitial absorption by grain boundaries in α -Fe. *Physical Review B*. 2012 Feb 10;85(6):064108.
 107. World Health Organization. Department of Mental Health, Substance Abuse, World Health Organization. Mental Health Evidence, Research Team. Mental health atlas 2005. World Health Organization; c2005.
 108. Shitaye JE, Tsegaye W, Pavlik I. Bovine tuberculosis infection in animal and human populations in Ethiopia: a review. *Veterinarni Medicina*. 2007;52(8):317-332.